

## Overweight and Obesity are Risk Factors of Severe Illness in Patients with COVID-19

Jian Wang<sup>1\*</sup>, Li Zhu<sup>2\*</sup>, Longgen Liu<sup>3\*</sup>, Xiang-an Zhao<sup>4\*</sup>, Zhaoping Zhang<sup>1\*</sup>, Leyang Xue<sup>5\*</sup>, Xuebing Yan<sup>6</sup>, Songping Huang<sup>7</sup>, Yang Li<sup>8</sup>, Juan Cheng<sup>9</sup>, Biao Zhang<sup>10</sup>, Tianmin Xu<sup>4</sup>, Chunyang Li<sup>6</sup>, Fang Ji<sup>6</sup>, Fang Ming<sup>7</sup>, Yun Zhao<sup>11</sup>, Huaping Shao<sup>12</sup>, Dawen Sang<sup>9</sup>, Haiyan Zhao<sup>12</sup>, Xinying Guan<sup>13</sup>, Xiaobing Chen<sup>14</sup>, Yuxin Chen<sup>15</sup>, Rahma Issa<sup>16</sup>, Jie Wei<sup>16</sup>, Rui Huang<sup>1</sup>, Chuanwu Zhu<sup>2</sup>, and Chao Wu<sup>1</sup>

**Objective:** This study aimed to observe the clinical characteristics of patients with coronavirus disease 2019 (COVID-19) with overweight and obesity.

**Methods:** Consecutive patients with COVID-19 from 10 hospitals of Jiangsu province, China, were enrolled.

**Results:** A total of 297 patients with COVID-19 were included, and 39.39% and 13.47% of patients had overweight and obesity, respectively. The proportions of bilateral pneumonia (92.50% vs. 73.57%,  $P=0.033$ ) and type 2 diabetes (17.50% vs. 3.57%,  $P=0.006$ ) were higher in patients with obesity than lean patients. The proportions of severe illness in patients with overweight (12.82% vs. 2.86%,  $P=0.006$ ) and obesity (25.00% vs. 2.86%,  $P<0.001$ ) were significantly higher than lean patients. More patients with obesity developed respiratory failure (20.00% vs. 2.86%,  $P<0.001$ ) and acute respiratory distress syndrome (5.00% vs. 0%,  $P=0.024$ ) than lean patients. The median days of hospitalization were longer in patients with obesity than lean patients (17.00 days vs. 14.00 days,  $P=0.029$ ). Overweight (OR, 4.222; 95% CI: 1.322-13.476;  $P=0.015$ ) and obesity (OR, 9.216; 95% CI: 2.581-32.903;  $P=0.001$ ) were independent risk factors of severe illness. Obesity (HR, 6.607; 95% CI: 1.955-22.329;  $P=0.002$ ) was an independent risk factor of respiratory failure.

**Conclusions:** Overweight and obesity were independent risk factors of severe illness in COVID-19 patients. More attention should be paid to these patients.

Obesity (2020) 28, 2049-2055.

### Introduction

Coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was declared a pandemic by the World Health Organization (1). Although most of the SARS-CoV-2 infection typically leads to relatively mild symptoms, 287,399 patients still died globally up to May 13, 2020 (2). Old age and comorbidities, such as hypertension, diabetes, and chronic respiratory disease, were identified as risk factors of poor outcomes for patients with COVID-19 according to previous studies (3,4). Obesity was regarded as a common risk factor to aggravate the severity of respiratory diseases, which was associated with poor prognosis in influenza A pulmonary

infection (5,6). Animal experiments have found that obesity alters inflammatory and pathological responses in the lung during influenza (7-9). Excessive adipose accumulation could result in insulin resistance, oxidative stress, chronic inflammation, and circulating nutrients abnormality (10,11). However, few studies have focused on the impacts of obesity on COVID-19.

A retrospective study that enrolled 124 patients with COVID-19 who were admitted to intensive care unit (ICU) showed more patients required invasive mechanical ventilation (IMV) therapy with increased BMI (12). Another study found that patients with COVID-19 who were aged <60 years with a BMI between 30 and 34 kg/m<sup>2</sup> had 2.0 and 1.8 times the

<sup>1</sup> Department of Infectious Diseases, Nanjing Drum Tower Hospital, The Affiliated Hospital of Nanjing University Medical School, Nanjing, China. Correspondence: Chao Wu (dr.wu@nju.edu.cn), Rui Huang (doctor\_hr@126.com), Chuanwu Zhu (zhuchw@126.com) <sup>2</sup> Department of Infectious Diseases, The Affiliated Infectious Diseases Hospital of Soochow University, Suzhou, China <sup>3</sup> Department of Infectious Diseases, The Third People's Hospital of Changzhou, Changzhou, China <sup>4</sup> Department of Gastroenterology, Northern Jiangsu People's Hospital, Clinical Medical College of Yangzhou University, Yangzhou, China <sup>5</sup> Department of Critical Medicine, Huai'an No. 4 People's Hospital, Huai'an, China <sup>6</sup> Department of Infectious Diseases, Affiliated Hospital of Xuzhou Medical University, Xuzhou, China <sup>7</sup> Department of Infectious Diseases, Nantong Third People's Hospital, Nantong University, Nantong, China <sup>8</sup> Department of Infectious Diseases, Taizhou People's Hospital, Taizhou, China <sup>9</sup> Department of Infectious Diseases, Yancheng Second People's Hospital, Yancheng, China <sup>10</sup> Department of Quality Control Office, Huai'an No. 4 People's Hospital, Huai'an, China <sup>11</sup> Department of Infectious Diseases, The Third People's Hospital of Yangzhou, Yangzhou, China <sup>12</sup> Department of Infectious Diseases, The People's Hospital of Suqian, Suqian, China <sup>13</sup> Department of Neurology, The Affiliated Hospital of Kangda College of Nanjing Medical University, The First People's Hospital of Lianyungang, Lianyungang, China <sup>14</sup> Department of Emergency, The Affiliated Hospital of Kangda College of Nanjing Medical University, The First People's Hospital of Lianyungang, Lianyungang, China <sup>15</sup> Department of Laboratory Medicine, Nanjing Drum Tower Hospital, The Affiliated Hospital of Nanjing University Medical School, Nanjing, China <sup>16</sup> Department of Infectious Diseases, Nanjing Drum Tower Hospital Clinical College of Nanjing Medical University, Nanjing, China.

\*Jian Wang, Li Zhu, Longgen Liu, Xiang-an Zhao, Zhaoping Zhang, and Leyang Xue contributed equally to this work.

© 2020 The Obesity Society. Received: 17 May 2020; Accepted: 20 July 2020; Published online 23 September 2020. doi:10.1002/oby.22979

risk for admission to acute and critical care as compared with individuals with a BMI <30 kg/m<sup>2</sup>, respectively (13). These studies suggested that obesity may be associated with the severity of COVID-19. However, for several confounders such as age, the presence of comorbidities were not adjusted, which might have impacted the results. Whether overweight and obesity are independent risk factors of severe COVID-19 requires further research. In addition, the sample sizes are relatively small in the previous studies (12,13). This study aimed to investigate the clinical features of patients with COVID-19 with overweight and obesity in a multicenter cohort of COVID-19 in Jiangsu province, China.

## Methods

### Study population

Between January 18, 2020, and February 26, 2020, 342 consecutive patients with COVID-19 from 10 medical centers in 10 cities of Jiangsu, China, were enrolled. All patients with COVID-19 were diagnosed by clinical manifestations, chest computed tomography, and real-time polymerase chain reaction according to World Health Organization interim guidance and the Guidelines for the Diagnosis and Treatment of Novel Coronavirus Infection by the National Health Commission (Trial Version 5) (14,15). All patients with COVID-19 were tested positive for SARS-CoV-2 by real-time polymerase chain reaction in throat swab specimens. The last followed-up date was February 29, 2020. The study was approved by the ethics review boards of these medical centers.

### Data collection and study definitions

We retrospectively recorded the clinical characteristics, complications, and outcomes of patients by electronic medical record system. The computational formula of BMI was weight (kilograms) divided by height (meters) squared. According to criterion of guidelines for prevention and control of overweight and obesity in Chinese adults,  $24 \leq \text{BMI} < 28$  and  $\text{BMI} \geq 28$  was defined as overweight and obesity, respectively (16,17). Severe COVID-19 was defined according to the current guideline as follows: (1) respiratory frequency  $\geq 30/\text{min}$ , (2) pulse oximeter oxygen saturation  $\leq 93\%$  at rest, and (3) oxygenation index  $\leq 300$  mmHg (15). Acute respiratory distress syndrome (ARDS) was defined according to the Berlin definition (18).

### Statistical analysis

Continuous variables were described as medians (interquartile range [IQR]), and categorical variables were presented as the counts and percentages. The independent group *t* tests (normal distribution) and Mann-Whitney U (non-normal distribution) were used to compare continuous variables between groups. Chi-square or Fisher exact test was used to compare the categorical variables. Multivariate logistic and cox regression analysis was used to adjust for confounding factors, including age, gender, and comorbidities.  $P < 0.05$  was considered to be statistically significant. SPSS version 22.0 software (SPSS Inc., Chicago, Illinois) was used for the analysis.

## Results

### Clinical characteristics of patients with overweight and obesity on admission

Thirty-four patients were excluded because of the lack of BMI data, and 11 patients under 12 years old were also excluded. Eventually,

297 patients were enrolled in this study. The clinical characteristics were presented in Table 1. Of the 297 patients with COVID-19, 117 (39.39%) and 40 (13.47%) had overweight ( $24 \leq \text{BMI} < 28$ ) and obesity ( $\text{BMI} \geq 28$ ), respectively. The median age was 38.00 (IQR, 31.25–54.00) years, 48.00 (IQR, 36.50–57.00) years, and 47.00 (IQR, 34.25–57.75) years in patients with lean ( $\text{BMI} < 24$ ), overweight, and obesity, respectively. Patients with overweight and obesity were older, and the proportion of male gender was higher in patients with overweight and obesity than lean patients. The median BMI was 22.17 (IQR, 20.55–23.12), 25.71 (IQR, 24.87–26.61), and 29.75 (IQR, 29.18–30.81) in these three groups, respectively. The proportions of a medical history of type 2 diabetes were significantly different in these three groups ( $P = 0.008$ ). More patients had a history of type 2 diabetes in the obesity group than the lean group (17.50% vs. 3.57%,  $P = 0.006$ ), whereas the proportion was comparable between the overweight group and lean group (11.11% vs. 3.57%,  $P = 0.054$ ). The clinical symptoms on admission were similar among the three groups. The median levels of fasting blood glucose (FBG) presented an increasing trend in patients who were lean and with overweight and obesity ( $P < 0.001$ ). Patients with obesity presented higher FBG levels than lean patients (6.48 mmol/L vs. 5.47 mmol/L,  $P < 0.001$ ), whereas patients with overweight had similar FBG levels with lean patients (5.69 mmol/L vs. 5.47 mmol/L,  $P = 0.119$ ). However, there were no significant differences in triglycerides and total cholesterol among the three groups. The proportion of bilateral pneumonia presented an increasing trend in patients who were lean and with overweight and obesity ( $P = 0.018$ ). More patients with obesity had bilateral pneumonia on admission than lean patients (92.50% vs. 73.57%,  $P = 0.033$ ), while there was no difference between patients with overweight and lean patients (82.91% vs. 73.57%,  $P = 0.219$ ).

### Treatment and clinical outcomes of patients with overweight and obesity

The proportions of patients who used atomized inhalation of interferon  $\alpha$ -2b (60.00%, 56.41%, and 57.50%,  $P = 0.84$ ), lopinavir-ritonavir (70.01%, 75.21%, and 80.00%,  $P = 0.383$ ), and arbidol (52.86%, 47.86%, and 52.50%,  $P = 0.709$ ) were comparable among the three groups (Table 2). The proportions of oxygen therapy ( $P = 0.009$ ) and non-IMV therapy ( $P = 0.003$ ) were significantly different in these three groups. More patients with overweight received oxygen therapy compared with lean patients (66.67% vs. 48.57%,  $P = 0.012$ ) during hospitalization, while the proportion of oxygen therapy was comparable between patients with obesity and lean patients (65.00% vs. 48.57%,  $P = 0.201$ ). Moreover, more patients with overweight (6.84% vs. 0.71,  $P = 0.024$ ) and obesity (12.50% vs. 0.71,  $P < 0.001$ ) received non-IMV therapy than lean patients. No patient received IMV treatment in our study. The proportions of patients who developed respiratory failure ( $P = 0.001$ ) and ARDS ( $P = 0.049$ ) were significantly different in these three groups. In total, 20% of patients with obesity developed respiratory failure, which was significantly higher than in lean patients (2.86%,  $P < 0.001$ ), while the proportion of respiratory failure was comparable between patients with overweight and lean patients (8.55% vs. 2.86%,  $P = 0.135$ ). Similarly, the proportions of patients developing ARDS were higher in patients with obesity than lean patients (5.00% vs. 0%,  $P = 0.024$ ), while the proportion of ARDS was comparable between patients with overweight and lean patients (1.71% vs. 0%,  $P = 0.360$ ). The proportions of severe illness were different in the three groups ( $P < 0.001$ ). More patients with overweight (12.82% vs. 2.86%,  $P = 0.006$ ) and obesity (25.00% vs. 2.86%,  $P < 0.001$ ) had severe illness than lean patients. However, the proportions of patients admitted to ICU were comparable among the three groups ( $P = 0.087$ ). The median days of hospitalization were different among the three groups ( $P = 0.025$ ). Patients

**TABLE 1** Demographic and clinical features of patients with COVID-19

Variables (n [%] or median [IQR])	Lean (n = 140)	Overweight (n = 117)	Obesity (n = 40)	P value
Age (y)	38.00 (31.25-54.00)	48.00 (36.50-57.00)	47.00 (34.25-57.75)	0.024
Male	68 (48.57)	72 (61.54)	24 (60.00)	0.092
BMI (kg/m <sup>2</sup> )	22.17 (20.55-23.12)	25.71 (24.87-26.61)	29.75 (29.18-30.81)	<0.001
<b>Comorbidities</b>				
Hypertension	15 (10.71)	24 (20.51)	9 (22.50)	0.053
Diabetes	5 (3.57)	13 (11.11)	7 (17.50)	0.008
Chronic lung diseases	4 (2.86)	6 (5.13)	2 (5.00)	0.620
Cardiovascular diseases	0	5 (4.27)	1 (2.50)	0.051
Malignant tumors	1 (0.71)	1 (0.85)	2 (5.00)	0.098
<b>Onset symptoms</b>				
Fever	89 (63.57)	84 (71.79)	32 (80.00)	0.099
Cough	75 (53.57)	70 (59.83)	23 (57.50)	0.597
Fatigue	24 (17.14)	24 (20.51)	13 (32.50)	0.106
Sore throat	16 (11.43)	12 (10.26)	5 (12.50)	0.914
Muscle ache	19 (13.57)	9 (7.69)	1 (2.50)	0.072
Headache	11 (7.86)	7 (5.98)	1 (2.50)	0.462
<b>Laboratory and imaging findings</b>				
WBC (×10 <sup>9</sup> /L)	4.46 (3.39-5.57)	5.10 (4.01-6.23)	5.23 (4.04-6.36)	0.006
Lymphocytes (×10 <sup>9</sup> /L)	1.18 (0.89-1.62)	1.23 (0.87-1.61)	1.11 (0.88-1.60)	0.879
ALT (U/L)	23.00 (15.00-30.00)	29.00 (22.00-40.00)	31.00 (23.00-48.50)	<0.001
Cr (μmol/L)	63.00 (52.50-77.15)	70.00 (57.90-82.00)	66.20 (59.05-82.55)	0.048
INR	1.05 (1.00-1.12)	1.04 (1.00-1.11)	1.02 (0.99-1.06)	0.284
UA (mmol/L)	263.75 (210.75-309.75)	272.20 (197.03-331.75)	295.90 (216.35-383.75)	0.482
FBG (mmol/L)	5.47 (4.90-6.11)	5.69 (5.08-6.50)	6.48 (5.45-7.89)	<0.001
TG (mmol/L)	1.09 (0.76-1.61)	1.27 (0.90-1.72)	1.37 (1.03-1.55)	0.063
TC (mmol/L)	3.75 (3.21-4.52)	3.83 (3.25-4.47)	3.80 (3.18-4.30)	0.968
<b>Chest CT</b>				
No pneumonia	14 (10.00)	5 (4.27)	2 (5.00)	0.175
Unilateral pneumonia	23 (16.43)	15 (12.82)	1 (2.50)	0.07
Bilateral pneumonia	103 (73.57)	97 (82.91)	37 (92.50)	0.018
Days from symptom onset to admission	5.00 (2.00-8.00)	6.00 (3.00-8.75)	4.50 (2.00-7.25)	0.069

ALT, alanine transaminase; Cr, creatinine; FBG, fasting blood glucose; INR, international normalized ratio; IQR, interquartile range; TC, total cholesterol; TG, triglyceride; UA, uric acid; WBC, white blood cells.

with obesity stayed longer in hospital than lean patients (17.00 days vs. 14.00 days,  $P=0.029$ ), while there was no significant difference of hospital stays between patients with overweight and lean patients (16.00 days vs. 14.00 days,  $P=0.421$ ).

The distributions of BMI categories were significantly different between patients with severe illness and nonsevere illness ( $P<0.001$ ), respiratory failure and nonrespiratory failure ( $P=0.001$ ), noninvasive mechanical ventilation and without noninvasive mechanical ventilation ( $P=0.003$ ) (Figure 1). However, although the proportions of patients with overweight and obesity had a higher rate of ICU admission, the difference was not statistically significant ( $P=0.087$ ).

### Associations of overweight and obesity with severe illness

Logistic regression analysis was performed to identify the association between obesity and severe illness (Table 3). Univariate

analysis presented that overweight (odds ratio [OR], 5.000; 95% CI: 1.611-15.516;  $P=0.005$ ), obesity (OR, 11.333; 95% CI: 3.329-38.583;  $P<0.001$ ), and type 2 diabetes (OR, 7.087; 95% CI: 2.782-18.053;  $P<0.001$ ) were associated with severe illness. In the multivariate analysis, overweight (OR, 4.222; 95% CI: 1.322-13.476;  $P=0.015$ ) and obesity (OR, 9.216; 95% CI: 2.581-32.903;  $P=0.001$ ) were independent risk factors of severe COVID-19 after adjusted age, sex, the presence of hypertension, and type 2 diabetes.


### Associations of overweight and obesity with respiratory failure

The associated factors of respiratory failure in patients with COVID-19 were analyzed by cox regression analysis (Table 4). The univariate analysis showed that the factors for respiratory failure were obesity (HR, 7.542; 95% CI: 2.270-25.055;  $P=0.001$ ) and type 2 diabetes (HR, 4.986; 95% CI: 1.949-12.756;  $P=0.001$ ). On multivariate analysis,

**TABLE 2** Treatment, complications, and clinical outcome of patients with COVID-19

Variables (n [%] or median [IQR])	Lean (n = 140)	Overweight (n = 117)	Obesity (n = 40)	P value
<b>Treatment</b>				
Atomized inhalation of interferon $\alpha$ -2b	84 (60.00)	66 (56.41)	23 (57.50)	0.84
Lopinavir-ritonavir	98 (70.00)	88 (75.21)	32 (80.00)	0.383
Arbidol	74 (52.86)	56 (47.86)	21 (52.50)	0.709
Oxygen therapy	68 (48.57)	78 (66.67)	26 (65.00)	0.009
Noninvasive mechanical ventilation	1 (0.71)	8 (6.84)	5 (12.50)	0.003
<b>Complications</b>				
Respiratory failure	4 (2.86)	10 (8.55)	8 (20.00)	0.001
ARDS	0	2 (1.71)	2 (5.00)	0.049
<b>Outcomes</b>				
Severe illness	4 (2.86)	15 (12.82)	10 (25.00)	<0.001
Admission to ICU	4 (2.86)	10 (8.55)	4 (10.00)	0.087
Days of Hospitalization	14.00 (12.00-19.00)	16.00 (12.00-20.00)	17.00 (14.00-22.50)	0.025

ARDS, acute respiratory distress syndrome; ICU, intensive care unit; IQR, interquartile range.



**Figure 1** Distributions of lean, overweight, and obesity of patients with COVID-19 with different conditions. ICU, intensive care unit; RF, respiratory failure; NMV, noninvasive mechanical ventilation.

obesity (HR, 6.607; 95% CI: 1.955-22.329;  $P=0.002$ ), type 2 diabetes (HR, 5.197; 95% CI: 1.837-14.699;  $P=0.002$ ), and age >60 years (HR, 2.766; 95% CI: 1.018-7.514;  $P=0.046$ ) were associated with an increased risk of respiratory failure.

## Discussion

Overweight and obesity are serious global health problems (19,20). The global prevalence ratios of overweight and obesity are 38.5% to

39.4% and 10.1% to 14.8%, respectively, in the general population (21). In our study, 39.39% and 13.47% of the patients with COVID-19 had overweight and obesity, respectively, suggesting that overweight and obesity may be not susceptible factors of COVID-19.

In our study, the most common symptoms were fever and cough, which were similar with previous studies (22,23). However, there were no significant differences in clinical symptoms among patients with different BMI. Patients with obesity had higher FBG levels and higher proportion of type 2 diabetes, indicating that obesity was associated with an increased risk for type 2 diabetes. Several studies have demonstrated that the presence of type 2 diabetes was a significant risk factor for severe illness and fatal outcome of COVID-19 (24-26). Thus, the association between the obesity-related comorbidities and severe COVID-19 deserves further investigation.

In our study, more patients with obesity had bilateral pneumonia compared with lean patients. In the report by Cai et al. (27), among 383 patients with COVID-19 from Shenzhen, China, obesity was also associated with a higher risk for severe pneumonia compared with lean patients. More patients with overweight and obesity received oxygen therapy and non-IMV in the present study. Our results revealed that overweight and obesity were risk factors of severe illness and more likely to develop complications such as respiratory failure and ARDS.

After adjusting the confounding factors such as age and sex, overweight and obesity were still independent risk factors of severe illness of COVID-19. Taken together, these results suggested that overweight and obesity were independently associated with the severity of COVID-19.

However, the mechanisms of overweight and obesity contributing to severe COVID-19 are not yet defined. Obesity has been regarded as a risk factor of severe illness and poor prognosis in many infectious diseases (28). Obesity induces systematically chronic inflammation by increasing the secretion of cytokines such as interleukin 6, interleukin 8, and tumor necrosis factor- $\alpha$ , which may aggravate the injury of lung parenchyma and bronchi (29-31). A previous study has also found that obesity might impair adaptive immune responses in influenza virus infection (32). A similar mechanism might exist in patients with COVID-19. In addition, obesity causes a decrease in protective cardiorespiratory reserve and immune dysfunction (33). Sattar et al. (33) also reported obesity could increase the risk of thrombosis, which is an unignorable risk factor of severe COVID-19. With regard to lung function, obesity reduces expiratory volume and forced vital capacity (33-35). In addition, animal models demonstrated that obesity leads to decreased natural killer cell cytotoxicity and increased mortality in influenza infection (36). However, the mechanisms of overweight and obesity in the severity of COVID-19 deserve further investigation.

TABLE 3 Logistic regression analysis of risk factors for patients with severe COVID-19

Variables	Univariate		Multivariate	
	OR (95% CI)	P	OR (95% CI)	P
<b>Age</b>				
≤60	Reference			
>60	1.577 (0.635-3.913)	0.326	1.410 (0.469-4.244)	0.541
<b>Sites</b>	1.023 (0.897-1.190)	0.770		
<b>Sex</b>				
Female	Reference			
Male	2.295 (0.982-5.363)	0.055	1.925 (0.729-5.088)	0.186
<b>BMI (kg/m<sup>2</sup>)</b>				
<24	Reference			
24-28	5.000 (1.611-15.516)	0.005	4.167 (1.296-13.402)	0.017
≥28	11.333 (3.329-38.583)	<0.001	9.028 (2.523-32.299)	0.001
<b>Hypertension</b>				
No	Reference			
Yes	1.090 (0.394-3.015)	0.868	0.435 (0.124-1.528)	0.194
<b>Type 2 diabetes</b>				
No	Reference			
Yes	7.087 (2.782-18.053)	<0.001	5.333 (1.800-15.800)	0.003
<b>Chronic lung diseases</b>				
No	Reference			
Yes	1.911 (0.398-9.178)	0.418	1.751 (0.322-9.511)	0.517
<b>Cardiovascular diseases</b>				
No	Reference			
Yes	1.879 (0.212-16.653)	0.571	0.968 (0.066-14.280)	0.981
<b>Malignant tumors</b>				
No	Reference			
Yes	3.155 (0.317-31.354)	0.327	1.768 (0.150-20.879)	0.651

OR, odds ratio.

TABLE 4 Cox regression analysis of risk factors for respiratory failure

Variables	Univariate		Multivariate	
	HR (95% CI)	P	HR (95% CI)	P
Age				
≤60	Reference			
>60	1.927 (0.754-4.926)	0.170	2.766 (1.018-7.514)	0.046
Sex				
Female	Reference			
Male	1.808 (0.737-4.434)	0.196	1.432 (0.537-3.824)	0.473
BMI (kg/m <sup>2</sup> )				
<24	Reference			
24-28	2.969 (0.931-9.467)	0.066	2.349 (0.708-7.798)	0.163
≥28	7.542 (2.270-25.055)	0.001	6.607 (1.955-22.329)	0.002
Hypertension				
No	Reference			
Yes	0.803 (0.238-2.713)	0.724	0.359 (0.092-1.400)	0.140
Type 2 diabetes				
No	Reference			
Yes	4.986 (1.949-12.756)	0.001	5.197 (1.837-14.699)	0.002
Chronic lung diseases				
No	Reference			
Yes	2.446 (0.572-10.467)	0.228	2.233 (0.489-10.198)	0.300
Cardiovascular diseases				
No	Reference			
Yes	0.048 (0.000-15003.613)	0.638		0.981
Malignant tumors				
No	Reference			
Yes	3.741 (0.503-27.835)	0.198	2.319 (0.285-18.878)	0.432

OR, odds ratio.

There were several limitations in our study. First, the outcomes of patients with COVID-19 had relatively favorable outcomes with no deaths. Thus, we could not analyze the association of overweight/obesity and fatal outcome in patients with COVID-19. Second, the associations of overweight/obesity with IMV could not be analyzed either. However, Kalligeros et al. (37) reported the potential association of obesity with severe outcomes in 102 patients hospitalized with COVID-19. They found that obesity was independently associated with the use of IMV (37). Third, many of the treatments and outcomes (oxygen therapy, non-IMV, admission to ICU, days of hospitalization) are subject to bias from clinicians who were in charge for the management of patients. Fourth, our study was conducted later in the pandemic, and by this point, people had suspicions and some studies were already conducted on the topic that overweight and obesity were risk factors for more severe COVID-19 outcomes. Thus, the clinicians in charge of care might have just been overly cautious, which might have biased our results. Furthermore, we could not include all the patients in our province. Thus, there is a potential selection bias in our study. However, nearly half of the confirmed cases in our province were included in our present study. We consider that our study is representative. Finally, the impacts of overweight and obesity on the long-term outcomes of patients with COVID-19 remain unclear.

In conclusion, patients with COVID-19 with overweight and obesity had higher risks for severe illness. Therefore, more attention should be

paid to patients with COVID-19 with overweight or obesity. However, more studies are needed to confirm our findings and to reveal the underlying mechanisms of overweight and obesity associated with higher risks for severe illness in COVID-19. **O**

**Funding agencies:** This study was supported by the Fundamental Research Funds for the Central Universities (No. 14380459).

**Disclosure:** The authors declared no conflict of interest exist.

**Author contributions:** Concept and design: CW, RH, and CZ; drafting of the manuscript: JWang, RH, and XZ; critical revision of the manuscript for important intellectual content: CW, RH, LX, LL, ZZ, FL, and CZ; statistical analysis: JWang; administrative, technical, or material support: LZ, XZ, ZZ, XY, SH, YL, JC, BZ, TX, CL, FJ, FM, YZ, HS, DS, HZ, XG, XC, and YC; supervision: CW; acquisition, analysis, or interpretation of data: JWei, LZ, LX, LL, XZ, ZZ, and RH. All authors reviewed and approved the final version.

## References

- World Health Organization. WHO characterizes COVID-19 as a pandemic. Rolling updates on coronavirus disease (COVID-19) website. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen>. Published March 11, 2020. Updated July 31, 2020. Accessed March 15, 2020.
- World Health Organization. Coronavirus disease 2019 (COVID-19) Situation Report – 55. [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200315-sitrep-55-covid-19.pdf?sfvrsn=33daa5cb\\_8](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200315-sitrep-55-covid-19.pdf?sfvrsn=33daa5cb_8). Published March 15, 2020. Accessed March 15, 2020.
- Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020;382:1708-1720.
- Chen T, Wu D, Chen H, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ* 2020;368:m1091. doi:10.1136/bmj.m1091

5. Milner JJ, Rebeles J, Dhungana S, et al. Obesity increases mortality and modulates the lung metabolome during pandemic H1N1 influenza virus infection in mice. *J Immunol* 2015;194:4846-4859.
6. Bijani B, Pahlevan AA, Qasemi-Barqi R, Jahanihashemi H. Metabolic syndrome as an independent risk factor of hypoxaemia in influenza A (H1N1) 2009 pandemic. *Infect Med* 2016;24:123-130.
7. O'Brien KB, Vogel P, Duan S, et al. Impaired wound healing predisposes obese mice to severe influenza virus infection. *J Infect Dis* 2012;205:252-261.
8. Karlsson EA, Sheridan PA, Beck MA. Diet-induced obesity impairs the T cell memory response to influenza virus infection. *J Immunol* 2010;184:3127-3133.
9. Milner JJ, Sheridan PA, Karlsson EA, Schultz-Cherry S, Shi Q, Beck MA. Diet-induced obese mice exhibit altered heterologous immunity during a secondary 2009 pandemic H1N1 infection. *J Immunol* 2013;191:2474-2485.
10. Milner JJ, Beck MA. The impact of obesity on the immune response to infection. *Proc Nutr Soc* 2012;71:298-306.
11. Johnson AR, Milner JJ, Makowski L. The inflammation highway: metabolism accelerates inflammatory traffic in obesity. *Immunol Rev* 2012;249:218-238.
12. Simonnet A, Chetboun M, Poissy J, et al. High prevalence of obesity in severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) requiring invasive mechanical ventilation. *Obesity (Silver Spring)* 2020;28:1195-1199.
13. Lighter J, Phillips M, Hochman S, et al. Obesity in patients younger than 60 years is a risk factor for Covid-19 hospital admission. *Clin Infect Dis* 2020;71:896-897.
14. World Health Organization. Clinical management of COVID-19. <https://www.who.int/publications/item/clinical-management-of-covid-19>. Accessed February 8, 2020.
15. National Health Commission of the People's Republic of China. Diagnosis and Treatment Protocol for COVID-19 (Trial Version 7). [http://en.nhc.gov.cn/2020-03/29/c\\_78469.htm](http://en.nhc.gov.cn/2020-03/29/c_78469.htm). Updated March 29, 2020. Accessed February 22, 2020.
16. Zhao S, Jia X, Fan X, et al. Association of obesity with the clinicopathological features of thyroid cancer in a large, operative population: a retrospective case-control study. *Medicine* 2019;98:e18213. doi:10.1097/MD.00000000000018213
17. Wang L, Du X, Dong JZ, et al. Body mass index and all-cause mortality in patients with atrial fibrillation: insights from the China atrial fibrillation registry study. *Clin Res Cardiol* 2019;108:1371-1380.
18. ARDS Definition Task Force; Ranieri VM, Rubenfeld GD, Thompson BT, et al. Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012;307:2526-2533.
19. Dehghan M, Akhtar-Danesh N, Merchant AT. Childhood obesity, prevalence and prevention. *Nutr J* 2005;4:24. doi:10.1186/1475-2891-4-24
20. Hossain P, Kawar B, El Nahas M. Obesity and diabetes in the developing world—a growing challenge. *N Engl J Med* 2007;356:213-215.
21. Chooi YC, Ding C, Magkos F. The epidemiology of obesity. *Metabolism* 2019;92:6-10.
22. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;395:507-513.
23. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497-506.
24. Yan Y, Yang Y, Wang F, et al. Clinical characteristics and outcomes of patients with severe covid-19 with diabetes. *BMJ Open Diabetes Res Care* 2020;8:e001343. doi:10.1136/bmjdc-2020-001343
25. Chen Q, Zheng Z, Zhang C, et al. Clinical characteristics of 145 patients with corona virus disease 2019 (COVID-19) in Taizhou, Zhejiang, China. *Infection* 2020;48:543-551. doi:10.1007/s15010-020-01432-5
26. Huang I, Lim MA, Pranata R. Diabetes mellitus is associated with increased mortality and severity of disease in COVID-19 pneumonia - a systematic review, meta-analysis, and meta-regression. *Diabetes Metab Syndr* 2020;14:395-403.
27. Cai Q, Chen F, Luo F, et al. Obesity and COVID-19 severity in a designated hospital in Shenzhen, China. *Diabetes Care* 2020;43:1392-1398.
28. Huttunen R, Syrjänen J. Obesity and the risk and outcome of infection. *Int J Obes (Lond)* 2013;37:333-340.
29. Zhang X, Zheng J, Zhang L, et al. Systemic inflammation mediates the detrimental effects of obesity on asthma control. *Allergy Asthma Proc* 2018;39:43-50.
30. Soto GJ, Frank AJ, Christiani DC, Gong MN. Body mass index and acute kidney injury in the acute respiratory distress syndrome. *Crit Care Med* 2012;40:2601-2608.
31. Bernejo-Martin JF, Ortiz de Lejarazu R, Pumarola T, et al. Th1 and Th17 hypercytokinemia as early host response signature in severe pandemic influenza. *Crit Care* 2009;13:R201. doi:10.1186/cc8208
32. Green WD, Beck MA. Obesity impairs the adaptive immune response to influenza virus. *Ann Am Thorac Soc* 2017;14(suppl 5):S406-S409.
33. Sattar N, McInnes IB, McMurray JJV. Obesity a risk factor for severe COVID-19 infection: multiple potential mechanisms. *Circulation* 2020;142:4-6.
34. Watson RA, Pride NB, Thomas EL, et al. Reduction of total lung capacity in obese men: comparison of total intrathoracic and gas volumes. *J Appl Physiol* 2010;108:1605-1612.
35. Jones RL, Nzekwu M-MU. The effects of body mass index on lung volumes. *Chest* 2006;130:827-833.
36. Smith AG, Sheridan PA, Harp JB, Beck MA. Diet-induced obese mice have increased mortality and altered immune responses when infected with influenza virus. *J Nutr* 2007;137:1236-1243.
37. Kalligeros M, Shehadeh F, Mylona EK, et al. Association of obesity with disease severity among patients with COVID-19. *Obesity (Silver Spring)* 2020;28:1200-1204.

CopyrightsPlatform